

## RESEARCH NOTE

### Is Vigor of Regeneration a Key Factor in Recovery from Peripheral Nerve Injuries?

LUIS DE MEDINACELI, TAM QUACH, ANNE-MARIE DUCHEMIN,  
AND RICHARD JED WYATT

Neuropsychiatry Branch, Intramural Research Program, National Institute of Mental Health,  
Saint Elizabeths Hospital, Washington, D.C. 20032

Received June 23, 1986

The nerve-growth-promoting effects of the tricyclic antidepressant, imipramine, were tested on the sympathetic ganglion of chickens and on the sciatic nerve of rats. A powerful neurotrophic action was observed *in vitro*, but the utilization of the drug *in vivo* did not modify the functional recovery from a crush lesion. © 1986 Academic Press, Inc.

Le Gros Clark's observation that nerve fiber regrowth is enhanced by salivary gland tissue (7) was followed by Bueker's work with mouse sarcoma tissue (1). Since then, nerve growth factor (NGF) has been used experimentally in both central and peripheral nervous systems. The *in vivo* effects of NGF, however, are difficult to analyze because it is quickly metabolized. Therefore, there is no answer to the following questions: (a) In order to obtain optimal enhancement of regeneration, should NGF be applied: to the cell body? to the neurite at the site of damage? or distally, with the idea of attracting and guiding the sprouts? (b) Would the increased vigor of regeneration thus produced be useful in clinical practice?

Recently, the tricyclic antidepressant, imipramine, was found by one of us (T.Q.) to have a strong NGF-like effect *in vitro*. Briefly, dissociated neurons from sympathetic ganglia of 12-day-old chick embryos were incubated 24 h in the absence (control), or in the presence of 1  $\mu$ M imipramine. The number of surviving neurons was determined by direct double-blind count. The

788

0014-4886/86 \$3.00

Copyright © 1986 by Academic Press, Inc.  
All rights of reproduction in any form reserved.

number of surviving neurons per milliliter was almost doubled in the presence of imipramine (Tam Quach, in preparation).

In the present study, we tested the behavioral effects of imipramine on the peripheral nervous system. Imipramine is relatively slowly catabolized and, when injected intraperitoneally, diffuses until even distribution is attained throughout the body. By insuring a long-term action of the drug all along the nervous tracts, this circumvented the first of the above questions.

Sprague-Dawley male rats weighing 200 to 250 g at the beginning of the experiment were anesthetized with pentobarbital and chloral hydrate, i.p. In all animals, the left sciatic nerve was aseptically exposed and crushed at mid thigh with maximum force for 1 min in the bare jaws of a serrated hemostat. The resulting lesion was 2 mm wide. In 19 animals, 2 mg/kg of imipramine were injected daily intraperitoneally; the treatment started 3 days before the procedure, and was continued until the 25th postoperative day. In another group of 18 rats, only solvent was injected.

The functional condition of the rats was assessed through the use of the sciatic functional index (SFI), based on measurements made from rats' walking tracks and expressed in units of functional deficit (3). Collection and analysis of data were made by a "blind" observer with a data management system (4). Animals were tested preoperatively and at regular postoperative intervals.




FIG. 1. Absence of effect of imipramine on the functional recovery from a crush lesion to the sciatic nerve of the rat. The mean ( $\pm$ SE) sciatic functional index expressed in units of deficit is shown as a function of time. The shaded area between +10 and -10 defines normality. Functional recovery was identical in both groups.

BEST AVAILABLE COPY

## RESEARCH NOTE

# Angiotensin II-Sensitive Neurons in the Rat Lateral Hypothalamic Area with Efferent Projections to the Subfornical Organ

J. TANAKA, H. KABA, H. SAITO, AND K. SETO<sup>1</sup>

*Department of Physiology, Kochi Medical School, Nankoku, Kochi 781-51, Japan*

*Received February 21, 1986; revision received July 7, 1986*

Thirteen neurons in the lateral hypothalamic area were antidromically activated by electrical stimulation of the subfornical organ in the rat. The activity of almost all identified neurons ( $N = 11$ ) was excited by microiontophoretically applied angiotensin II. On the other hand, of the antidromically unidentified lateral hypothalamic area neurons ( $N = 10$ ) tested, 4 were excited by microiontophoretically applied angiotensin II and 6 were not affected. The excitatory responses of identified and unidentified lateral hypothalamic area neurons to angiotensin II were blocked by microiontophoretically applied saralasin, an angiotensin II antagonist. © 1986 Academic Press, Inc.

The subfornical organ (SFO) is implicated in angiotensin II (AII)-induced drinking (1, 7, 10, 14), a centrally mediated pressor response (4, 7, 10), and the control of vasopressin release from the neurohypophysis (3, 6). Recent immunohistochemical studies have identified AII-immunoreactive pathways from the lateral hypothalamic area (LHA) to the SFO (8, 9), suggesting the importance of these pathways for the control of body fluid balance and central cardiovascular regulation.

In a previous study (5), we described that LHA efferent fibers have an excitatory influence on the activity of SFO neurons projecting to the supraoptic nucleus. Because the LHA receives the AII-immunoreactive effer-

Abbreviations: LHA—lateral hypothalamic area, SFO—subfornical organ, MIPh—microiontophoretically, AII—angiotensin II, Sar—saralasin.

<sup>1</sup> We thank Conrad Zagory, Jr. for helpful advice in the preparation of this manuscript. The work was supported in part by grant 59770126 from the Ministry of Education, Science and Culture, Japan. Please address correspondence to Dr. Tanaka.

791

0014-4886/86 \$3.00

Copyright © 1986 by Academic Press, Inc.  
All rights of reproduction in any form reserved.

No difference was found in functional recovery of the two groups (Fig. 1); in all animals, recovery commenced at about day 15 and was satisfactory by day 21.

The results obtained for the period of functional recovery (days 11 through 25) were submitted to a one-way multiple analysis of variance (MANOVA). The test was not significant from day 11 [ $F(1,36) = 0.47, P = 0.49$ ] through day 25 ( $F = 0.00, P = 0.96$ ), confirming that imipramine had no clinically detectable effects on sciatic nerve regrowth.

Three interpretations could be offered for the lack of efficacy of imipramine *in vivo*: (a) Imipramine may act exclusively on sensory neurons. Evaluation of sensory levels in animals is notoriously imprecise (2), and our method of functional assessment could be insufficient to demonstrate the beneficial effects of the drug. The general reliability of this method, however, has been independently confirmed (5, 6, 8). (b) The effects of increased vigor of regeneration might be difficult to measure in the rat because of the short distance the sprouts have to grow before reaching their target. Moderate and short-lived differences could thus be overlooked. Over greater distance, i.e., in longer nerves, differences would be magnified, last longer and become measurable. (c) Vigor of regeneration might not be a key factor in the recovery from peripheral nerve injuries. We believe that this last interpretation is correct; the unpredictable functional results of peripheral nerve surgery can be explained without reference to growth promoting substances (L. de Medinaceli, in preparation).

## REFERENCES

1. BUEKER, E. D. 1948. Implantation of tumors in the hindlimb field of the chick and the developmental response of the lumbosacral nervous system. *Anat. Rec.* 102: 369-390.
2. DEEKE, L., AND C. H. TATOR. 1973. Neurophysiological assessment of afferent and efferent conduction in the injured spinal cord of monkeys. *J. Neurosurg.* 39: 65-74.
3. DE MEDINACELI, L., W. J. FREED, AND R. J. WYATT. 1982. An index of the functional condition of rat sciatic nerve based on measurements made from walking tracks. *Exp. Neurol.* 77: 634-643.
4. DE MEDINACELI, L., E. DERENZO, AND R. J. WYATT. 1984. Rat sciatic functional index data management system with digitized input. *Comput. Biomed. Res.* 17: 185-192.
5. GOLDBERG, N. H., S. S. DESHPANDE, C. S. MAY, F. K. TOY, R. T. SMOOT, C. K. MESHUL, AND E. X. ALBUQUERQUE. 1984. Disparity between neurophysiologic measurements and clinical reality following peripheral nerve transection and microneurography. *Surg. Forum* 35: 608-610.
6. KERNS, J. M., A. J. FAKHOURI, I. M. PAVKOVIC, AND K. J. SMITH. 1986. A twitch tension method to assess motor nerve function. *J. Neurosci. Methods*, in press.
7. LE GROS CLARK, W. E. 1942. The problem of neuronal regeneration in the central nervous system. II. The insertion of peripheral nerve stumps into the brain. *J. Anat.* 77: 251-259.
8. TERZIS, J. K., AND K. J. SMITH. 1986. "de Medinaceli" versus microsuturing; a critical appraisal of nerve repair. *Proc. Am. Soc. Reconstr. Microsurg.* 2: 116.